

# Math 32B - Fall 2019

## Practice Exam 1

Full Name: Solutions

UID: \_\_\_\_\_

Circle the name of your TA and the day of your discussion:

Steven Gagniere

Jason Snyder

Ryan Wilkinson

Tuesday

Thursday



### Instructions:

- Read each problem carefully.
- Show all work clearly and circle or box your final answer where appropriate.
- Justify your answers. A correct final answer without valid reasoning will not receive credit.
- Simplify your answers as much as possible.
- Include units with your answer where applicable.
- Calculators are not allowed but you may have a  $3 \times 5$  inch notecard.

---

Page	Points	Score
1	20	
2	20	
3	15	
4	20	
5	25	
Total:	100	

1. (10 points) Evaluate the iterated integral.




no easy antiderivative

both vertically simple  
and horizontally simple

$$\begin{aligned} \int_0^4 \int_{\sqrt{y}}^2 \sqrt{x^3+1} \, dx \, dy &= \int_0^2 \int_0^{x^2} \sqrt{x^3+1} \, dy \, dx = \int_0^2 [y\sqrt{x^3+1}]_0^{x^2} \, dx \\ &= \int_0^2 x^2 \sqrt{x^3+1} \, dx = \left[ \frac{1}{2} \cdot \frac{2}{3} (x^3+1)^{3/2} \right]_0^2 = \frac{2}{9} [(x^3+1)^{3/2}]_0^2 \\ &\quad \text{Should be } 1/3 \\ &= \frac{2}{9} (9^{3/2} - 1^{3/2}) = \frac{2}{9} (27 - 1) = \frac{2 \cdot 26}{9} = \boxed{\frac{52}{9}} \end{aligned}$$

2. (10 points) Evaluate the iterated integral.




$$\begin{aligned} x &= r \cos \theta \\ y &= r \sin \theta \\ x^2 + y^2 &= r^2 \end{aligned}$$

polar


$$\int_0^3 \int_0^{\sqrt{9-x^2}} e^{x^2+y^2} \, dy \, dx \quad \text{no elementary antiderivative}$$

$$\begin{aligned} \int_0^3 \int_0^{\sqrt{9-x^2}} e^{x^2+y^2} \, dy \, dx &= \int_0^{\pi/2} \int_0^3 e^{r^2} r \, dr \, d\theta \\ &= \int_0^{\pi/2} d\theta \cdot \int_0^3 r e^{r^2} \, dr \\ &= [\theta]_0^{\pi/2} \cdot \left[ \frac{1}{2} e^{r^2} \right]_0^3 \\ &= \frac{\pi}{2} \cdot \frac{1}{2} (e^9 - e^0) = \boxed{\frac{\pi}{4} (e^9 - 1)} \end{aligned}$$

3. (10 points) Find the volume of the solid enclosed by  $z = 0$ ,  $y = z$ , and  $x^2 + y^2 = 4$ .




4. (10 points) Use a triple integral to find the volume of the solid enclosed by  $y = x^2$ ,  $z = 3y$ , and  $z = 2 + y$ .




5. (15 points) Consider the tetrahedron bounded by the coordinate planes and the plane  $x + y + z = 1$  with density function  $\delta(x, y, z) = 12y$ .

$$z = 1 - x - y$$

1. Find the mass of the tetrahedron.



$$\text{if } z=0, x+y=1 \text{ so } y=1-x$$



$$m = \iiint_W \delta \, dV = \int_0^1 \int_0^{1-x} \int_0^{1-x-y} 12y \, dz \, dy \, dx$$

$$\begin{aligned} &= \int_0^1 \int_0^{1-x} [12yz]_0^{1-x-y} \, dy \, dx = \int_0^1 \int_0^{1-x} 12y(1-x-y) \, dy \, dx \\ &= \int_0^1 \int_0^{1-x} 12y(1-x) - 12y^2 \, dy \, dx = \int_0^1 [6y^2(1-x) - 4y^3]_0^{1-x} \, dx \\ &= \int_0^1 6(1-x)^3 - 4(1-x)^3 \, dx = \int_0^1 2(1-x)^3 \, dx \\ &= \left[ -\frac{2}{4}(1-x)^4 \right]_0^1 = 0 + \frac{1}{2} = \boxed{\frac{1}{2}} \end{aligned}$$

(also x-simple  
and y-simple)

2. Set up but DO NOT EVALUATE the integrals used to find the center of mass of the tetrahedron.

center of mass is  $(x_{cm}, y_{cm}, z_{cm})$  with

$$x_{cm} = \frac{1}{m} \iiint_W x \delta(x, y, z) \, dV = 2 \int_0^1 \int_0^{1-x} \int_0^{1-x-y} 12xy \, dz \, dy \, dx$$

$$y_{cm} = \frac{1}{m} \iiint_W y \delta(x, y, z) \, dV = 2 \int_0^1 \int_0^{1-x} \int_0^{1-x-y} 12y^2 \, dz \, dy \, dx$$

$$z_{cm} = \frac{1}{m} \iiint_W z \delta(x, y, z) \, dV = 2 \int_0^1 \int_0^{1-x} \int_0^{1-x-y} 12yz \, dz \, dy \, dx$$

6. (20 points) Evaluate the triple integral  $\iiint_E x^2 dV$  where  $E$  is the solid above  $z = 0$  and inside  $4x^2 + 9y^2 + z^2 = 36$ .

Ellipsoid  $2x=u, 3y=v, z=w \quad \begin{matrix} u \\ v \\ w \end{matrix}$   
 $x=\frac{u}{2}, y=\frac{v}{3}, z=w \quad \begin{matrix} u^2+v^2+w^2=36 \\ u \\ v \end{matrix}$   
 Sphere  $\rho=6$   
 above  $w=0$

$$J(G) = \frac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{3} & 0 \\ 0 & 0 & 1 \end{vmatrix} = \frac{1}{6} \quad |J| = \frac{1}{6}$$

$$\text{so } \iiint_E x^2 dV = \iiint_W (\frac{u}{2})^2 \cdot \frac{1}{6} du dv dw = \frac{1}{24} \iiint_W u^2 du dv dw$$

Now spherical coordinates

$$u = \rho \sin\phi \cos\theta, \quad v = \rho \sin\phi \sin\theta, \quad w = \rho \cos\phi$$

$$0 \leq \rho \leq 6$$

$$0 \leq \theta \leq 2\pi$$

$$0 \leq \phi \leq \frac{\pi}{2}$$

$$\iiint_E x^2 dV = \frac{1}{24} \iiint_W u^2 du dv dw = \frac{1}{24} \int_0^{\pi/2} \int_0^{2\pi} \int_0^6 \rho^2 \sin^2\phi \cos^2\theta \cdot \rho^2 \sin\phi \, d\rho d\theta d\phi$$

$$= \frac{1}{24} \int_0^{\pi/2} \int_0^{2\pi} \int_0^6 \rho^4 \sin^3\phi \cos^2\theta \, d\rho d\theta d\phi$$

$$= \frac{1}{24} \int_0^{\pi/2} \sin^3\phi \, d\phi \cdot \int_0^{2\pi} \cos^2\theta \, d\theta \int_0^6 \rho^4 \, d\rho$$

$$= \frac{1}{24} \int_0^{\pi/2} \sin\phi (1 - \cos^2\phi) \, d\phi \cdot \int_0^{2\pi} \frac{1}{2}(1 + \cos(2\theta)) \, d\theta \int_0^6 \rho^4 \, d\rho$$


$$= \frac{1}{24} \cdot \frac{1}{2} \int_0^{\pi/2} \sin\phi - \sin\phi \cos^2\phi \, d\phi \cdot \int_0^{2\pi} 1 + \cos(2\theta) \, d\theta \int_0^6 \rho^4 \, d\rho$$

$$= \frac{1}{24} \cdot \frac{1}{2} \left[ -\cos\phi + \frac{\cos^3\phi}{3} \right]_0^{\pi/2} \cdot \left[ \theta + \frac{1}{2}\sin(2\theta) \right]_0^{2\pi} \cdot \left[ \frac{\rho^5}{5} \right]_0^6$$

$$= \frac{1}{24} \cdot \frac{1}{2} \left( 0 + 0 + 1 - \frac{1}{3} \right) (2\pi + 0 - 0 - 0) \left( \frac{6^5}{5} \right)$$

$$= \frac{1}{48} \cdot \frac{1}{2} \cdot \frac{\pi}{3} \cdot 2\pi \cdot \frac{6^5}{5} = \frac{6^3 \pi}{5} = \boxed{\frac{216\pi}{5}}$$

7. (10 points) Find the area inside one petal of the polar rose  $r = \sin(2\theta)$ .




$$0 \leq r \leq \sin(2\theta)$$

$$0 \leq \theta \leq \frac{\pi}{2}$$


$$\begin{aligned} A &= \iint_D 1 \, dA = \int_0^{\pi/2} \int_0^{\sin(2\theta)} 1 \cdot r \, dr \, d\theta \\ &= \int_0^{\pi/2} \left[ \frac{r^2}{2} \right]_0^{\sin(2\theta)} d\theta = \int_0^{\pi/2} \frac{\sin^2(2\theta)}{2} d\theta \\ &= \int_0^{\pi/2} \frac{1}{2} \left( 1 - \cos(4\theta) \right) d\theta = \frac{1}{4} \int_0^{\pi/2} 1 - \cos(4\theta) d\theta \\ &= \frac{1}{4} \left[ \theta - \frac{1}{4} \sin(4\theta) \right]_0^{\pi/2} = \frac{1}{4} (\pi/2 - 0 - 0 + 0) \\ &= \boxed{\frac{\pi}{8}} \end{aligned}$$

8. (15 points) Use a change of variables to evaluate  $\iint_D x \, dA$  where  $D$  is the region in the first quadrant bounded by  $y = 0$ ,  $y = 4$ ,  $y = x^2$ , and  $y = x^2 - 4$ .



horizontally simple but required to use a change of variables

$$0 \leq y \leq 4, \quad 0 \leq x^2 - y \leq 4$$



so let  $u = y$ ,  $v = x^2 - y$  then  $0 \leq u \leq 4$ ,  $0 \leq v \leq 4$

$$J(G^{-1}) = \frac{\partial(u, v)}{\partial(x, y)} = \begin{vmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{vmatrix} \stackrel{G^{-1}}{=} \begin{vmatrix} 0 & 1 \\ 2x & -1 \end{vmatrix} = -2x \text{ in region } D \quad x \geq 0$$

$$J(G) = \frac{1}{|J(G^{-1})|} = -\frac{1}{2x}, \quad |J(G)| = \frac{1}{2x}. \quad (\text{can solve } x = \sqrt{u+v} \text{ but not necessary})$$

$$\iint_D x \, dA = \iint_R x \cdot \frac{1}{2x} \, du \, dv = \iint_R \frac{1}{2} \, du \, dv = \frac{1}{2} \text{Area}(R) = \frac{1}{2} (4^2) = \boxed{8}$$

$$\text{check: } \int_0^4 \int_{\sqrt{y+4}}^{\sqrt{y+4}} x \, dx \, dy = \int_0^4 \left[ \frac{x^2}{2} \right]_{\sqrt{y+4}}^{\sqrt{y+4}} dy = \frac{1}{2} \int_0^4 y+4 - y \, dy = \frac{1}{2} \int_0^4 4 \, dy = 8$$